



XVII Международная астрономическая олимпиада
XVII International Astronomy Olympiad

Корея, Кванджу

16 – 24. X. 2012

Gwangju, Korea

язык	<i>English</i>
language	

Theoretical round. Problems to solve

General note. Maybe not all problems have correct questions. Some questions (maybe the main question of the problem, maybe one of the subquestions) may make no real sense. In this case you have to write in your answer (in English or Russian): «impossible situation – ситуация невозможна». Of course, this answer has to be explained numerically or logically.

Data from the tables (Planetary data, stars, constants, etc.) may be used for solving every problem.

The answers «Да-Yes» or «Нет-No» have to be written in English or Russian.

- 1. Transit of Venus.** Recently, on June 6, 2012, an infrequent astronomical phenomenon, transit of Venus across the solar disc, took place. The next transit of Venus will take place only in 2117. Calculate the date of that transit. (Answer without calculations will not be considered even as a partial solution.)
- 2. Transit of Pseudovenus.** Recently, on June 6, 2012, an infrequent astronomical phenomenon, transit of Venus across the solar disc took place. Suppose somebody did not understand the phenomenon and ascribed it not to transit of real Venus but of some moon, which we name Pseudovenus, rotating around the Earth in a circular orbit. Find the radius of the orbit of Pseudovenus and diameter of this sky body. Effects due to axial rotating of the Earth should not be taken into account.
- 3. Old persons' star.** There is ancient legend in Korea that says, if you managed to see the “Old persons' star” thrice, you are lucky person and will live a long life. The “Old persons' star”, now known as Canopus, was seen brighter and better in past times, but even now sometimes one can see this star in Korea. Estimate approximately what visible stellar magnitude Canopus may have when observing it from the southern coast of Jeju island (Korea) in the most favorable conditions. The territory of the island is located at latitudes between $33^{\circ}12' \text{ N}$ and $33^{\circ}34' \text{ N}$ and longitudes between $126^{\circ}09' \text{ E}$ and $126^{\circ}57' \text{ E}$. Take from the tables and recollect for yourself the necessary additional information.
- 4. Stars on Mars.** As you know, last year the Polar Bear (whom you have already met in the texts of many International Astronomy Olympiads) arrived to Mars for astronomical observations. Nowadays his friend Penguin also made a fascinating journey to Mars. At the same instant of time, the Bear and the Penguin observe stars in zenith and see Canopus and Sirius respectively. Estimate roughly, what is the distance (measured on the Martian surface) between the animals? At what height above the horizon does the Bear observe Sirius? The solution has to include a picture with an image of the Bear and the Penguin on Mars. Necessary sizes or angular sizes should be in the picture. Recollect for yourself the necessary information about the Polar Bear and Penguin.
- 5. Venus and Earth.** At what maximum distance from the Venus ecliptic the Earth can be visible in the sky from Venus (actually, from a point outside the Venus atmosphere)? Orbits of the planets may be considered circular.
- 6. Parallaxes.** In our part of the Galaxy the mean distance between the stars is about 6 light years. Assume that an interferometer can measure parallaxes with an error of ± 0.001 arc second. How many stars of our Galaxy could have their parallax determined by this interferometer?



XVII Международная астрономическая олимпиада
XVII International Astronomy Olympiad

Корея, Кванджу

16 – 24. X. 2012

Gwangju, Korea

язык	<u>Русский</u>
language	
язык	<u>English</u>
language	

Элементы орбит.

Физические характеристики некоторых планет, Луны, Солнца и Эриды

Parameters of orbits.

Physical characteristics of some planets, Moon, Sun and Eris

Небесное тело, планета	Среднее расстояние от центрального тела		Сидерический (или аналогичный) период обращения		На- клон орби- ты, <i>i</i>	Экс- цен- три- ситет, <i>e</i>	Эквато- риальн. диаметр <i>км</i>	Масса <i>10²⁴ кг</i>	Сред- няя плот- ность <i>г/см³</i>	Ускор. своб. пад. у пов. <i>м/с²</i>	На- клон оси	Макс. блеск, вид. с Земли (**)	Аль- бедо
	в <i>астр. ед.</i>	в <i>млн. км</i>	в тропич. годах	в средних сутках									
Body, planet	Average distance to central body		Sidereal period (or analogous)		Orbital inclination, <i>i</i>	Ec- centri- city <i>e</i>	Equat. diameter <i>km</i>	Mass <i>10²⁴ kg</i>	Av. den- sity <i>g/cm³</i>	Grav. accelr. at surf. <i>m/s²</i>	Axial tilt	Max. magn. From Earth (**)	Al- bedo
	in <i>astr. units</i>	in <i>10⁶ km</i>	in tropical years	in days									
Солнце Sun	$1,6 \cdot 10^9$	$2,5 \cdot 10^{11}$	$2,2 \cdot 10^8$	$8 \cdot 10^{10}$			1392000	1989000	1,409			-26,74 ^m	
Меркурий Mercury	0,387	57,9	0,241	87,969	7,00°	0,206	4 879	0,3302	5,43	3,70	0,01°		0,06
Венера Venus	0,723	108,2	0,615	224,7007	3,40	0,007	12 104	4,8690	5,24	8,87	177,36		0,78
Земля Earth	1,000	149,6	1,000	365,2564	0,00	0,017	12 756	5,9742	5,515	9,81	23,44		0,36
Луна Moon	0,00257	0,38440	0,0748	27,3217	5,15	0,055	3 475	0,0735	3,34	1,62	6,7	-12,7 ^m	0,07
Марс Mars	1,524	227,9	1,880	686,98	1,85	0,093	6 794	0,6419	3,94	3,71	25,19	-2,0 ^m	0,15
Юпитер Jupiter	5,204	778,6	11,862	4 332,59	1,30	0,048	142 984	1899,8	1,33	24,86	3,13	-2,7 ^m	0,66
Сатурн Saturn	9,584	1433,7	29,458	10 759,20	2,48	0,054	120 536	568,50	0,70	10,41	26,73	0,7 ^m	0,68
Эрида Eris	68,05			205 029	43,82	0,435	2 326	0,0167	2,52	0,7			0,96

**) Для Луны – в среднем противостоянии.

**) For Moon – in mean opposition.



XVII Международная астрономическая олимпиада
XVII International Astronomy Olympiad

Корея, Кванджу

16 – 24. X. 2012

Gwangju, Korea

язык	<u>Русский</u>
language	
язык	<u>English</u>
language	

Некоторые константы и формулы

Some constants and formulae

Скорость света в вакууме, c (м/с)	299 792 458	Speed of light in vacuum, c (m/s)
Гравитационная постоянная, G ($\text{Н} \cdot \text{м}^2/\text{кг}^2$)	$6.674 \cdot 10^{-11}$	Constant of gravitation, G ($\text{Н} \cdot \text{м}^2/\text{кг}^2$)
Солнечная постоянная, A ($\text{Вт}/\text{м}^2$)	1367	Solar constant, A (W/m^2)
Параметр Хаббла, среднее значение	71	mean value Hubble parameter,
H_0 (км/с/Мпк) диапазон значений	50-100	diapason of values H_0 (km/s/Mpc)
Постоянная Планка, h (Дж·с)	$6.626 \cdot 10^{-34}$	Plank constant, h (J·s)
Заряд электрона, e (Кл)	$1.602 \cdot 10^{-19}$	Charge of electron, e (C)
Масса электрона, m_e (кг)	$9.109 \cdot 10^{-31}$	Mass of electron, m_e (kg)
Соотношение масс протона и электрона	1836.15	Proton-to-electron ratio
Постоянная Фарадея, F (Кл/моль)	96 485	Faraday constant, F (C/mol)
Магнитная постоянная, μ_0 (Гн/м)	$1.257 \cdot 10^{-6}$	Magnetic constant, μ_0 (H/m)
Универсальная газовая постоянная, R (Дж/моль/К)	8.314	Universal gas constant, R (J/mol/K)
Постоянная Больцмана, k (Дж/К)	$1.381 \cdot 10^{-23}$	Boltzmann constant, k (J/K)
Постоянная Стефана-Больцмана, σ ($\text{Вт}/\text{м}^2/\text{K}^4$)	$5.670 \cdot 10^{-8}$	Stefan-Boltzmann constant, σ ($\text{W}/\text{m}^2/\text{K}^4$)
Константа смещения Вина, b (м·К)	0.002897	Wien's displacement constant, b (m·K)
Лабораторная длина волны $\text{H}\alpha$ (Å)	6562.81	Laboratory wavelength of $\text{H}\alpha$ (Å)
Длина тропического года, T (сут)	365.242199	Tropical year length, T (days)
Стандартная атмосфера (Па)	101 325	Standard atmosphere (Pa)
Ослабление видимого света земной атмосферой в зените (минимально)	19%, 0.23^m	Visible light extinction by the terrestrial atmosphere in zenith (minimum)
Показатель преломления воды при 20°C, n	1.334	Refractive index of water for 20°C, n
Момент инерции шара	$I = \frac{2}{5} MR^2$	Moment of inertia of a solid ball
Площадь сферы	$S = 4\pi R^2$	Area of sphere
π	3.14159265	π
e	2.71828183	e
Золотое сечение, ϕ	1.61803399	Golden ratio, ϕ



язык	<u>Русский</u>
language	
язык	<u>English</u>
language	

XVII Международная астрономическая олимпиада
XVII International Astronomy Olympiad

Корея, Кванджу

16 – 24. X. 2012

Gwangju, Korea

Данные о некоторых звёздах

Data of some stars

			RA	DEC	ρ	m	SC	масса mass
Солнце	Sun	$\odot$	$0^h - 24^h$	$-23^\circ 26' + 23^\circ 26'$	$8''.794$	$-26^m.74$	G2	$1 M_\odot$
Альдебаран	Aldebaran	α Tau	$04^h 35^m 55^s$	$16^\circ 30' 33''$	$0''.048$	$0^m.85^v$	K5	$2.5 M_\odot$
Альтаир	Altair	α Aql	$19^h 50^m 47^s$	$08^\circ 52' 06''$	$0''.195$	$0^m.77$	A7	$1.7 M_\odot$
Антарес	Antares	α Sco	$16^h 29^m 24^s$	$-26^\circ 25' 55''$	$0''.006$	$0^m.96$	M1+B4	$22.4 M_\odot$
Арктур	Arcturus	α Boo	$14^h 15^m 40^s$	$19^\circ 10' 57''$	$0''.089$	$-0^m.04^v$	K1	$1.1 M_\odot$
Ахернар	Achernar	α Eri	$01^h 37^m 43^s$	$-57^\circ 14' 12''$	$0''.026$	$0^m.46$	B3	
зв.Барнарда	Barnard's star	Oph	$17^h 57^m 48^s$	$04^\circ 41' 36''$	$0''.545$	$9^m.54$	M4	
Бетельгейзе	Betelgeuse	α Ori	$05^h 55^m 10^s$	$07^\circ 24' 25''$	$0''.005$	$0^m.5^v$	M1	
Вега	Vega	α Lyr	$18^h 36^m 56^s$	$38^\circ 47' 01''$	$0''.129$	$0^m.03$	A0	
Денеб	Deneb	α Cyg	$20^h 41^m 26^s$	$45^\circ 16' 49''$	$0''.002$	$1^m.25$	A2	
Канопус	Canopus	α Car	$06^h 23^m 57^s$	$-52^\circ 41' 45''$	$0''.010$	$-0^m.72$	F0	
Капелла	Capella	α Aur	$05^h 16^m 41^s$	$45^\circ 59' 53''$	$0''.073$	$0^m.08$	G5+G0	
Полярная	Polaris	α UMi	$02^h 31^m 49^s$	$89^\circ 15' 51''$	$0''.0076$	$1^m.97^v$	F7	
Процион	Procyon	α CMi	$07^h 39^m 18^s$	$05^\circ 13' 30''$	$0''.288$	$0^m.38$	F5	
Ригель	Rigel	β Ori	$05^h 14^m 32^s$	$-08^\circ 12' 06''$	$0''.013$	$0^m.12$	B8	
Сириус	Sirius	α CMa	$06^h 45^m 09^s$	$-16^\circ 42' 58''$	$0''.375$	$-1^m.46$	A1	
Спика	Spica	α Vir	$13^h 25^m 12^s$	$-11^\circ 09' 41''$	$0''.023$	$0^m.98$	B1	
Альфа Центавра	Alpha Centauri	α Cen	$14^h 39^m 36^s$	$-60^\circ 50' 07''$	$0''.751$	$-0^m.01$ $1^m.33$	G2 K1	$2.0 M_\odot$
Бета Центавра	Beta Centauri	β Cen	$14^h 03^m 49^s$	$-60^\circ 22' 23''$	$0''.009$	$0^m.61$	B1	$21 M_\odot$
Эпсилон Эридана	Epsilon Eridani	ϵ Eri	$03^h 32^m 56^s$	$-09^\circ 27' 30''$	$0''.311$	$3^m.74$	K2	$0.82 M_\odot$