

**[Solution]***Theoretical Question 3**Thermal Vibration of Surface Atoms*

(1) (a) The wavelength of the incident electron is

$$\begin{aligned}
 \lambda &= \frac{h}{p} = \frac{h}{\sqrt{2meV}} \\
 &= \frac{6.63 \times 10^{-34}}{\sqrt{2 \times 9.11 \times 10^{-31} \times 1.60 \times 10^{-19} \times 64.0}} \\
 &= 1.53 \times 10^{-10} \text{ m} = 1.53 \text{ \AA}
 \end{aligned}$$

(b) Consider the interference between the atomic rows on the surface as shown in Fig. 3c.




Fig.3c

The path difference between electron beam 1 and 2 is

$$\Delta \ell = b(\sin \phi - \sin \phi_0) = n\lambda$$

Given  $\phi_0 = 15.0^\circ$ ,  $\lambda = 1.53 \text{ \AA}$  and  $b = \frac{a}{\sqrt{2}} = \frac{3.92}{\sqrt{2}} = 2.77 \text{ \AA}$ , two solutions are possible.

(i) When  $n = 0$ ,  $\phi = \phi_0 = 15.0^\circ$

(Answer 1)

(ii) When  $n = 1$

$$\Delta \ell = 2.77(\sin \phi - \sin 15^\circ) = 1 \times 1.53$$

$$\sin \phi = \frac{1.53 + 0.72}{2.77} = 0.812$$

**[Solution]** (continued) Theoretical Question 3


### Vibration of Surface Atoms

$$\phi = 54.3^\circ \quad (\text{Answer 2})$$

For  $n = 2$ , no solution exists as  $\Delta\ell = 2.77(\sin\phi - \sin 15^\circ) = 2 \times 1.53$  and  $\sin\phi > 1$ .

$$(2) \quad I = I_0 \exp\langle -(\vec{u} \cdot \Delta\vec{K})^2 \rangle$$

Fig. 3d



For the specularly reflected beam, we have from Fig. 3d

$$\Delta\vec{K} = \vec{K}' - \vec{K} = 2K \cos\theta \hat{x}$$

where  $\hat{x}$  is the unit vector in the direction of the surface normal. Take the  $x$ -component of  $\vec{u}$ , we then obtain

$$I = I_0 e^{-\langle u_x^2(t) \cdot 4K^2 \cos^2\theta \rangle} = I_0 e^{-4K^2 \cos^2\theta \langle u_x^2(t) \rangle} \quad (2)$$

The vibration in the direction of the surface normal of the surface atoms is simple harmonic, take

$$u_x(t) = A \cos \omega t$$

$$\text{Q} \quad \langle u_x^2(t) \rangle = \frac{1}{\tau} \int_0^\tau u^2 dt = \frac{1}{\tau} \int_0^\tau A^2 \cos^2 \omega t dt = \frac{A^2}{\tau} \cdot \frac{\tau}{2} = \frac{A^2}{2}$$

$$\therefore A^2 = 2\langle u_x^2(t) \rangle$$

The total energy  $E$  is thus given by

$$E = \frac{1}{2} C A^2 = \frac{1}{2} C \cdot 2 \langle u_x^2(t) \rangle = C \langle u_x^2(t) \rangle = m' \omega^2 \langle u_x^2(t) \rangle$$

**[Solution]** (continued)      *Theoretical Question 3*  
*Vibration of Surface Atoms*

Therefore, one obtains

$$\langle u_x^2(t) \rangle = E / (m' \omega^2)$$

$$E = m' \omega^2 \langle u_x^2 \rangle = k_B T$$

where  $m'$  is the mass of the atom. From either of the above two equations, one then has the following equality

$$\langle u_x^2 \rangle = \frac{k_B T}{m' \omega^2} = \frac{k_B T}{m' 4\pi^2 f^2} \quad (3)$$

From eq. (3) and eq. (2), one obtains

$$I = I_0 e^{-4K^2 \cos^2 \theta \frac{k_B T}{m' 4\pi^2 f^2}}$$

where  $K = \frac{2\pi p}{h} = \frac{2\pi}{\lambda}$ . Accordingly,

$$I = I_0 e^{-\frac{4k_B \cos^2 \theta}{m' f^2 \lambda^2} T} = I_0 e^{-M' T} \quad (4)$$

and

$$\ln \frac{I}{I_0} = -M' T$$

From the plot of  $\ln \frac{I}{I_0}$  versus  $T$ , one obtains the slope

$$M' = \frac{4k_B \cos^2 \theta}{m' f^2 \lambda^2} \quad (5)$$

The slope of the curve can be estimated from Fig. 3b and leads to the result

$$M' = 2.3 \times 10^{-3}.$$

Using the following data in Eq.(5),

$$k_B = 1.38 \times 10^{-23} \text{ J / K}$$

$$\lambda = 1.53 \times 10^{-10} \text{ m}$$

$$m' = 195.1 \times 10^{-3} / (6.02 \times 10^{23}) = 3.24 \times 10^{-25} \text{ kg/atom}$$

**[Solution]** (continued)      *Theoretical Question 3*

*Vibration of Surface Atoms*

one finds

$$2.3 \times 10^{-3} = \frac{4 \times 1.38 \times 10^{-23} \cdot \cos^2 15^\circ}{\frac{195.1 \times 10^{-3}}{6.02 \times 10^{23}} \times f^2 \times (1.53 \times 10^{-10})^2}$$

The solution for frequency is then

$$f^2 = 3.0 \times 10^{24} \text{ (new)} \Rightarrow f = 1.7 \times 10^{12} \text{ Hz} \quad \text{Answer (a)}$$

From  $\langle u_x^2 \rangle = \frac{k_B T}{m' 4\pi^2 f^2}$ ,  $T = 300 \text{ K}$ , one finally obtains

$$\langle u_x^2 \rangle = \frac{1.38 \times 10^{-23} \cdot 300}{\frac{195.1 \times 10^{-3}}{6.02 \times 10^{23}} \times 4\pi^2 \times 3.0 \times 10^{24}} = 1.1 \times 10^{-22} \text{ m}^2 \text{ (new)}$$

and

$$\sqrt{\langle u_x^2 \rangle} = 1.0 \times 10^{-11} \text{ m} = 0.10 \text{ \AA} \text{ (new)}$$

Ans