

- (a) **(2 points)** Taking the center of the circular cross section of the filament as the origin O , find the electric potential at any point (x,z) very near the filament in terms of V_a , a and b where V_a is the electric potential of the surface of the filament, a is the radius of the filament and b is the distance between the center of the filament and the cylindrical wire mesh. (Ignore mirror charges.)
- (b) **(4 points)** An incoming electron plane wave with wave vector k_z is deflected by the “biprism” due to the x -component of the force exerted on the electron. Determine k_x the x -component of the wave vector due to the “biprism” in terms of the electron charge, e , v_z , V_a , k_z , a and b , where e and v_z are the charge and the z -component of the velocity of the electrons ($k_x \ll k_z$). Note that $\vec{k} = \frac{2\pi\vec{p}}{h}$ where h is the Planck constant.
- (c) Before the point S , electrons are emitted from a field emission tip and accelerated through a potential V_0 . Determine the wavelength of the electron in terms of the (rest) mass m , charge e and V_0 ,
- (i) **(2 points)** assuming relativistic effects can be ignored, and
- (ii) **(3 points)** taking relativistic effects into consideration.

- (d) In Tonomura *et al* experiment,

$$v_z = c/2,$$

$$V_a = 10 \text{ V},$$

$$V_0 = 50 \text{ kV},$$

$$a = 0.5 \text{ }\mu\text{m},$$

$$b = 5 \text{ mm},$$

$$\ell = 25 \text{ cm},$$

$$L = 1.5 \text{ m},$$

$$h = 6.6 \times 10^{-34} \text{ Js},$$

$$\text{electron charge, } e = 1.6 \times 10^{-19} \text{ C},$$

$$\text{mass of electron, } m_0 = 9.1 \times 10^{-31} \text{ kg},$$

$$\text{and the speed of light in vacuo, } c = 3 \times 10^8 \text{ ms}^{-1}$$

- (i) **(2 points)** calculate the value of k_x ,
- (ii) **(2 points)** determine the fringe separation of the interference pattern on the screen,
- (iii) **(1 point)** If the electron wave is a spherical wave instead of a plane wave, is the fringe spacing larger, the same or smaller than the fringe spacing calculated in (ii)?
- (iv) **(2 points)** In part (c), determine the percentage error in the wavelength of the electron using non-relativistic approximation.
- (v) **(2 points)** the distance d between the apparent double slits.